

KNOT FLOER HOMOLOGY AND SEIFERT SURFACES

ANDRÁS JUHÁSZ

ABSTRACT. Let K be a knot in S^3 of genus g and let $n > 0$. We show that if $\text{rk } \widehat{HFK}(K, g) < 2^{n+1}$ (where $\widehat{HFK}$ denotes knot Floer homology), in particular if K is an alternating knot such that the leading coefficient a_g of its Alexander polynomial satisfies $|a_g| < 2^{n+1}$, then K has at most n pairwise disjoint non-isotopic genus g Seifert surfaces. For $n = 1$ this implies that K has a unique minimal genus Seifert surface up to isotopy.

1. INTRODUCTION AND PRELIMINARIES

If S_1 and S_2 are Seifert surfaces of a knot $K \subset S^3$ then S_1 and S_2 are said to be equivalent if $S_1 \cap X(K)$ and $S_2 \cap X(K)$ are ambient isotopic in the knot exterior $X(K) = S^3 \setminus N(K)$, where $N(K)$ is a regular neighborhood of K . In [4] Kakimizu assigned a simplicial complex $MS(K)$ to every knot K in S^3 as follows.

Definition 1.1. $MS(K)$ is a simplicial complex whose vertices are the equivalence classes of the minimal genus Seifert surfaces of K . The equivalence classes $\sigma_0, \dots, \sigma_n$ span an n -simplex if and only if for each $0 \leq i \leq n$ there is a representative S_i of σ_i such that the surfaces $S_0, \dots, S_n$ are pairwise disjoint.

In [10] it is shown that the complex $MS(K)$ is always connected. I.e., if S and T are minimal genus Seifert surfaces for a knot K then there is a sequence $S = S_1, S_2, \dots, S_k = T$ of minimal genus Seifert surfaces such that $S_i \cap S_{i+1} = \emptyset$ for $0 \leq i \leq k-1$.

The main goal of this short note is to show that for a genus g knot K and for $n > 0$ the condition $\text{rk } \widehat{HFK}(K, g) < 2^{n+1}$ implies $\dim MS(K) < n$, consequently for $n = 1$ the knot K has a unique Seifert surface up to equivalence. This condition involves the use of knot Floer homology introduced by Ozsváth and Szabó in [8] and independently by Rasmussen in [9]. However, when K is alternating then this condition is equivalent to $|a_g| < 2^{n+1}$, where a_g is the leading coefficient of the Alexander polynomial of K . The alternating case is already a new result whose statement doesn't involve knot Floer homology. On the other hand, the proof of this particular case seems to need sutured Floer homology techniques, which is a generalization of knot Floer homology that was introduced by the author in [3]. At the time of writing this paper there are very few results in knot theory which can only be proved using Floer homology methods.

Date: February 2, 2008.

1991 Mathematics Subject Classification. 57M27; 57R58.

Key words and phrases. Alexander polynomial; Seifert surface; Floer homology.

Research partially supported by OTKA grant no. T49449.

The above statement does not hold for $n = 0$ since every knot has at least one minimal genus Seifert surface. However, it was shown in [6] and [2] that $\widehat{\text{rk}} \widehat{HFK}(K, g) < 2$ implies that the knot K is fibred.

To a knot K in S^3 and every $j \in \mathbb{Z}$ knot Floer homology assigns a graded abelian group $\widehat{HFK}(K, j)$ whose Euler characteristic is the coefficient a_j of the Alexander polynomial $\Delta_K(t)$. In [7] it is shown that if K is alternating then $\widehat{HFK}(K, j)$ is non-zero in at most one grading, thus $\widehat{\text{rk}} \widehat{HFK}(K, j) = |a_j|$.

Next we are going to review some necessary definitions and results from the theory of sutured manifolds and sutured Floer homology. Sutured manifolds were introduced by Gabai in [1].

Definition 1.2. A *sutured manifold* (M, γ) is a compact oriented 3-manifold M with boundary together with a set $\gamma \subset \partial M$ of pairwise disjoint annuli $A(\gamma)$ and tori $T(\gamma)$. Furthermore, the interior of each component of $A(\gamma)$ contains a *suture*, i.e., a homologically nontrivial oriented simple closed curve. We denote the union of the sutures by $s(\gamma)$.

Finally every component of $R(\gamma) = \partial M \setminus \text{Int}(\gamma)$ is oriented. Define $R_+(\gamma)$ (or $R_-(\gamma)$) to be those components of $\partial M \setminus \text{Int}(\gamma)$ whose normal vectors point out of (into) M . The orientation on $R(\gamma)$ must be coherent with respect to $s(\gamma)$, i.e., if δ is a component of $\partial R(\gamma)$ and is given the boundary orientation, then δ must represent the same homology class in $H_1(\gamma)$ as some suture.

A sutured manifold is called *taut* if $R(\gamma)$ is incompressible and Thurston norm minimizing in $H_2(M, \gamma)$.

The following definition was introduced in [3].

Definition 1.3. A sutured manifold (M, γ) is called *balanced* if M has no closed components, $\chi(R_+(\gamma)) = \chi(R_-(\gamma))$, and the map $\pi_0(A(\gamma)) \rightarrow \pi_0(\partial M)$ is surjective.

Example 1.4. If R is a Seifert surface of a knot K in S^3 then we can associate to it a balanced sutured manifold $S^3(R) = (M, \gamma)$ such that $M = S^3 \setminus (R \times I)$ and $\gamma = K \times I$. Observe that $R_-(\gamma) = R \times \{0\}$ and $R_+(\gamma) = R \times \{1\}$. Furthermore, $S^3(R)$ is taut if and only if R is of minimal genus.

Sutured Floer homology is an invariant of balanced sutured manifolds defined by the author in [3], and is a common generalization of the invariants $\widehat{HF}$ and $\widehat{HFK}$. It assigns an abelian group $SFH(M, \gamma)$ to each balanced sutured manifold (M, γ) . The following theorem is a special case of [2, Theorem 1.5].

Theorem 1.5. *Let K be a genus g knot in S^3 and suppose that R is a minimal genus Seifert surface for K . Then*

$$SFH(S^3(R)) \approx \widehat{HFK}(K, g).$$

A sutured manifold (M, γ) is called a *product* if it is homeomorphic to $(\Sigma \times I, \partial \Sigma \times I)$, where Σ is an oriented surface with boundary. If (M, γ) is a product then $SFH(M, \gamma) \approx \mathbb{Z}$. Let us recall [2, Theorem 1.4] and [2, Theorem 9.3].

Theorem 1.6. *If (M, γ) is a taut balanced sutured manifold then $SFH(M, \gamma) \geq \mathbb{Z}$. Furthermore, if (M, γ) is not a product then $SFH(M, \gamma) \geq \mathbb{Z}^2$.*

Definition 1.7. Let (M, γ) be a balanced sutured manifold. An oriented surface $S \subset M$ is called a *horizontal surface* if S is open, $\partial S = s(\gamma)$ in an oriented sense; moreover, $[S] = [R_+(\gamma)]$ in $H_2(M, \gamma)$, and $\chi(S) = \chi(R_+(\gamma))$.

A horizontal surface S defines a horizontal decomposition

$$(M, \gamma) \rightsquigarrow^S (M_-, \gamma_-) \coprod (M_+, \gamma_+)$$

as follows. Let $M_{\pm}$ be the union of the components of $M \setminus \text{Int}(N(S))$ that intersect $R_{\pm}(\gamma)$. Similarly, let $\gamma_{\pm}$ be the union of the components of $\gamma \setminus \text{Int}(N(S))$ that intersect $R_{\pm}(\gamma)$.

The following proposition is a special case of [2, Proposition 8.6].

Proposition 1.8. *Suppose that (M, γ) is a taut balanced sutured manifold and let S be a horizontal surface in it. Then*

$$\text{rk SFH}(M, \gamma) = \text{rk SFH}(M_-, \gamma_-) \cdot \text{rk SFH}(M_+, \gamma_+).$$

The following definition can be found for example in [6].

Definition 1.9. A balanced sutured manifold (M, γ) is called *horizontally prime* if every horizontal surface S in (M, γ) is isotopic to either $R_+(\gamma)$ or $R_-(\gamma)$ rel γ .

2. THE RESULTS

Theorem 2.1. *Let (M, γ) be a taut balanced sutured manifold such that both $R_+(\gamma)$ and $R_-(\gamma)$ are connected. Suppose that there is a sequence of pairwise disjoint non-isotopic connected horizontal surfaces $R_-(\gamma) = S_0, S_1, \dots, S_n = R_+(\gamma)$. Then*

$$\text{rk SFH}(M, \gamma) \geq 2^n.$$

Proof. We prove the theorem using induction on n . If $n = 1$ then (M, γ) is not a product since $R_-(\gamma)$ and $R_+(\gamma)$ are non-isotopic. Thus Theorem 1.6 implies that $\text{rk SFH}(M, \gamma) \geq 2$.

Now suppose that the theorem is true for $n - 1$. Since each S_k is connected we can suppose without loss of generality that S_1 separates S_i and S_0 for every $i \geq 2$. Let (M_-, γ_-) and (M_+, γ_+) be the sutured manifolds obtained after horizontally decomposing (M, γ) along S_1 . Note that both (M_-, γ_-) and (M_+, γ_+) are taut. Since S_0 and S_1 are non-isotopic (M_-, γ_-) is not a product so as before $\text{rk SFH}(M_-, \gamma_-) \geq 2$. Applying the induction hypothesis to (M_+, γ_+) and to the surfaces $R_-(\gamma_+), S_2, \dots, S_n = R_+(\gamma_+)$ we get that $\text{rk SFH}(M_+, \gamma_+) \geq 2^{n-1}$. So using Proposition 1.8 we see that $\text{rk SFH}(M, \gamma) \geq 2^n$. $\square$

Corollary 2.2. *If (M, γ) is a taut balanced sutured manifold and $\text{rk SFH}(M, \gamma) < 4$ then (M, γ) is horizontally prime. More generally, if $n > 0$ and $\text{rk SFH}(M, \gamma) < 2^{n+1}$ then (M, γ) can be cut into horizontally prime pieces by less than n horizontal decompositions.*

Proof. Suppose that $\text{rk SFH}(M, \gamma) < 2^{n+1}$. If (M, γ) is not horizontally prime then there is a surface S_1 in (M, γ) which is not isotopic to $R_{\pm}(\gamma)$. Decomposing (M, γ) along S_1 we get two sutured manifolds (M_-, γ_-) and (M_+, γ_+) . If they are not both horizontally prime then repeat the above process with a non-prime piece and obtain a horizontal surface S_2 , etc. This process has to end in less than n steps according to Theorem 2.1. $\square$

Theorem 2.3. *Let K be a knot in S^3 of genus g and let $n > 0$. If $\widehat{\text{rk HFK}}(K, g) < 2^{n+1}$ then K has at most n pairwise disjoint non-isotopic genus g Seifert surfaces, in other words, $\dim MS(K) < n$. If $n = 1$ then K has a unique Seifert surface up to equivalence.*

Proof. Suppose that $R, S_1, \dots, S_n$ are pairwise disjoint non-isotopic Seifert surfaces for K . According to Theorem 1.5 we have $\widehat{HFK}(K, g) \approx SFH(S^3(R))$. Let $S^3(R) = (M, \gamma)$. If $R_+(\gamma)$ and $R_-(\gamma)$ were isotopic then (M, γ) would be a product and S_1 and R would be equivalent. So the surfaces $R_-(\gamma) = S_0, S_1, \dots, S_n, S_{n+1} = R_+(\gamma)$ satisfy the conditions of Theorem 2.1, thus $\text{rk } SFH(S^3(R)) \geq 2^{n+1}$, a contradiction.

In particular, if $n = 1$ then $\dim MS(K) = 0$. But according to [10] the complex $MS(K)$ is connected, so it consists of a single point. $\square$

Corollary 2.4. *Suppose that K is an alternating knot in S^3 of genus g and let $n > 0$. If the leading coefficient a_g of its Alexander polynomial satisfies $|a_g| < 2^{n+1}$ then $\dim MS(K) < n$. If $|a_g| < 4$ then K has a unique Seifert surface up to equivalence.*

Proof. This follows from Theorem 2.3 and the fact that for alternating knots $\text{rk } \widehat{HFK}(K, g) = |a_g|$. $\square$

Remark 2.5. In [5] Kakimizu classified the minimal genus Seifert surfaces of all the prime knots with at most 10 crossings. The $n = 1$ case of Corollary 2.4 is sharp since the knot 7_4 is alternating, the leading coefficient of its Alexander polynomial is 4, and has 2 inequivalent minimal genus Seifert surfaces. On the other hand, the Alexander polynomial of the alternating knot 9_2 is also 4, but has a unique minimal genus Seifert surface up to equivalence.

Also note that [2, Theorem 1.7] implies that if the leading coefficient a_g of the Alexander polynomial of an alternating knot K satisfies $|a_g| < 4$ then the knot exterior $X(K)$ admits a depth ≤ 1 taut foliation transversal to $\partial X(K)$. Indeed, for alternating knots $g = g(K)$ and $|a_g| = \text{rk } \widehat{HFK}(K, g) \neq 0$, so the conditions of [2, Theorem 1.7] are satisfied.

ACKNOWLEDGEMENT

I would like to thank Zoltán Szabó for his interest in this work and Ko Honda and Tamás Kálmán for inspiring conversations.

REFERENCES

1. D. Gabai, *Foliations and the topology of 3-manifolds*, J. Differential Geom. **18** (1983), 445–503.
2. A. Juhász, *Floer homology and surface decompositions*, math.GT/0609779.
3. ———, *Holomorphic discs and sutured manifolds*, Algebr. Geom. Topol. **6** (2006), 1429–1457.
4. O. Kakimizu, *Finding disjoint incompressible spanning surfaces for a link*, Hiroshima Math. J. **22** (1992), no. 2.
5. ———, *Classification of the incompressible spanning surfaces for prime knots of 10 or less crossings*, Hiroshima Math. J. **35** (2005), no. 1.
6. Y. Ni, *Knot Floer homology detects fibred knots*, math.GT/0607156.
7. P. Ozsváth and Z. Szabó, *Heegaard Floer homology and alternating knots*, Geom. Topol. **7** (2003), 225–254.
8. ———, *Holomorphic disks and knot invariants*, Adv. Math. **186** (2004), no. 1, 58–116.
9. J. A. Rasmussen, *Floer homology and knot complements*, PhD thesis, Harvard University (2003).
10. M. Scharlemann and A. Thompson, *Finding disjoint Seifert surfaces*, Bull. London Math. Soc. **20** (1988), no. 1.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544, USA
E-mail address: `ajuhasz@math.princeton.edu`